

9600184

THE UNITED STATES OF AMERICA

South Ankota Agricultural Experiment Station

MILECORS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED, PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT. IN SE UNITED STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS A CLASS OF SEED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS SPECIFIED BY THE OWNER OF

IS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

WHEAT

'Russ'

In Testimonn Marrest, I have hereunto set my hand and caused the seal of the Hunt Anriety Protection Office to be affixed at the City of Washington, D.C. this twenty-eighth day of June in the year of our Lord one thousand nine hundred and ninety-six.

Allest:

Marsha A. Hamfor-Commissioner Plant Variety Protection Office

Secretary of Agriculture

10

EXHIBIT A. Russ (SD8073) Origin and Breeding History of the Variety

Russ is an F₆ derived line from the cross 'SD8052/SD2971' made at Brookings, South Dakota in 1984. The pedigree of SD8052 is (ND585, Butte*3/ (ND507, Waldron/5/ (RL4205, Pembina*6/2/ Thatcher*3/ Transfer/4/ Pembina*6/3/ Thatcher*2/2/ Marquis*6/ Red Egyptian)))/ Shield and the pedigree of SD2971 is Agent/3/ND441//Waldron/Bluebird/4/Butte/5/Len. The F₁ plants were grown at Weslaco, Texas during the winter of 1984-1985. Individual F₂ plant selections were made at Brookings, South Dakota in 1986 and were grown at Weslaco the following winter as plant rows. The plant rows at Weslaco were harvested as rows and used to plant F_{2:4} yield trials and a space planted nursery at Brookings in 1987. Based on data collected from the yield trials, individual plants were selected within the selected populations. Populations were selected based on grain yield, grain volume weight, and bread-making characteristics and individual plants were visually selected for resistance to prevalent foliar pathogens (viz. leaf rust and stem rust). Plant rows were grown in Weslaco during the winter and $F_{4:6}$ yield trials and space planted nursery were conducted at Brookings in 1988. As in 1987, the yield trial data was used to identify high yielding populations and individual plants were selected within the selected populations in the space planted nursery. The F₆ plant rows were grown in Weslaco and a single row was harvested and given the designation SD8073. Seed increase was conducted by the South Dakota Spring Wheat Breeding Program from 1989 through 1992. Breeders' seed was produced in 1993 and Foundation seed was produced in 1994.

Russ was tested by the South Dakota Spring Wheat Breeding Program from 1989 through 1994 and in the Uniform Regional Spring Wheat Nursery from 1991 through 1993. SD8073 was in the Spring Wheat Crop Quality Test in 1993 and 1994.

Russ has been uniform and stable for all morphological characters during the past four generations of selfing and increase. A tall variant (10 cm taller) was identified in the breeders seed at a frequency of approximately 0.2%. Up to 0.5% variant plants may be encountered in subsequent generations.

EXHIBIT B. Russ (SD8073) Statement of Distinctness

Russ is most similar to the hard red spring wheat cultivars 'Butte 86' and 'Sharp', but differs in the following characteristics:

- Russ is two days later to heading than Butte 86 and Sharp (Table 1.)
- Russ has a longer dough mixing time and a stronger mix pattern than Butte 86 and Sharp (Table 2.)
- Russ is heterogeneous for resistance to Hessian Fly, whereas, Butte 86 and Sharp do not have Hessian Fly resistance (see letter from Dr. J.H. Hatchett).

deleted per Cetter of MAY7,188 MAH 5-15-91

Table 1. South Dakota State University, Spring Wheat Breeding Trials Combined Over Locations.

| | | 19 | 93 | |
|------------|---|-------------------------------|-------------------------------|---------------------|
| | Yield bushels/acre (8) ¹ | Test Weight pounds/bushel (8) | Heading day of year (5) | Height cm (8) |
| Russ | 39.1 | 54.2 | 177.8 | 91.6 |
| Butte 86 | 37.9 | 55.4 | 175.1 | 89.3 |
| Sharp | 36.5 | 57.3 | 175.3 | 91.0 |
| Prospect | 35.0 | 52.6 | 179.3 | 88.3 |
| 2375 | 40.0 | 57.2 | 177.2 | 87.3 |
| LSD (0.05) | 3.81 | 1.27 | 0.96 | 3.69 |
| C.V. % | 10.78 | 2.33 | 0.43 | 4.19 |

| | | 19 | 94 | |
|------------|---|-------------------------------|-------------------------------|---------------------|
| | Yield bushels/acre (9) ¹ | Test Weight pounds/bushel (9) | Heading day of year (7) | Height cm (6) |
| Russ | 41.7 | 58.9 | 172.7 | 82.0 |
| Butte 86 | 38.3 | 58.6 | 171.4 | 83.3 |
| Sharp | 39.6 | 58.9 | 170.7 | 84.3 |
| Prospect | 36.7 | 57.7 | 174.8 | 79.5 |
| 2375 | 37.3 | 59.2 | 173.2 | 76.3 |
| LSD (0.05) | 2.76 | 1.28 | 1.09 | 3.27 |
| C.V. % | 7.91 | 2.36 | 0.60 | 4.19 |

¹ number of locations that data was collected.

Table 2. Quality characteristics. Samples from the 1992-1994 SDSU Advanced Yield Trial were evaluated by the Spring Wheat Quality Laboratory, Fargo, ND

| • | | Weig | | P | roteir %- | · · | Н | ardne | SS | | | action |
|----------|-----------|-----------|-----------|------------------|--------------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| | 92 (6) | 93 (5) | 94 (5) | 92 (6) | 93 (5) | 94 (5) | 92 (6) | 93 (5) | 94 (5) | 92 (6) | 93 (5) | 94 (5) |
| | (0) | (0) | (0) | (0) | (0) | (0) | (0) | (0) | (0) | (0) | (0) | (0) |
| Russ | 59.1 | 56.3 | 61.3 | 13.6 | 14.1 | 14.1 | 73 | 68 | 83 | 64.6 | 65.0 | 69.4 |
| Butte 86 | 59,6 | 57.6 | 61.8 | 13.8 | 14.1 | 14.4 | 81 | 75 | 83 | 66.7 | 66.0 | 69.5 |
| 2375 | 59.8 | 58.6 | 61.4 | 13.6 | 13.7 | 14.1 | 70 | 66 | 71 | 66.6 | 66.0 | 69.3 |
| Prospect | 58.7 | 55.9 | 61.3 | 13.2 | 13.5 | 13.2 | 56 | 49 | 58 | 61.5 | 63.7 | 68.3 |
| Sharp | 60.8 | 59.3 | 63.1 | 13.7 | 13.9 | 14.1 | 62 | 53 | 60 | 64.7 | 63.5 | 69.3 |
| | | | | , | | | | | | ٠ | | |
| | Mixi | ng Pat | ttern | Mix [*] | Time | (min) | Loa | f Vol | ıme | | | |
| | 92 | 93 | 94 | 92 | 93 | `94´ | 92 | 93 | 94 | | | |
| ****** | (6) | (5) | (5) | (6) | (5) | (5) | (6) | (5) | (5) | | | |
| Russ | 4.7 | 3.6 | 4.0 | 5.0 | 5.5 | 4.6 | 183 | 193 | 182 | | | - |
| Butte 86 | 3.3 | 2.2 | 2.0 | 3.1 | 3.5 | 3.4 | 183 | 197 | 179 | | | |
| 2375 | 3.3 | 2.4 | 3.0 | 3.5 | 4.0 | 3.9 | 185 | 194 | 182 | | | |
| Prospect | 1.7 | 1.4 | 2.0 | 2.9 | 3.4 | 3.3 | 183 | 190 | 180 | | | |
| Sharp | 3.5 | 2.4 | 2.0 | 3.3 | 3.6 | 3.2 | 184 | 196 | 182 | | | |

Photograph 1. South Dakota Spring Wheat Acid PAGE, Seed Testing Laboratory, South Dakota State University.




Table 3.

Seedling reaction of entries of the 1993 Uniform Hard Red Spring Wheat Performance Nursery to selected isolates of <u>Puccinia graminis</u> f. sp. <u>tritici</u>. (D.V. McVey, USDA-ARS, Cereal Rust Laboratory, U. of Minnesota, St. Paul, MN. 55108)

| | | | | | oduced | | olates | | | |
|-----|------------|------------|-------|------|--------|----------------|--------|------|-------------|-------------|
| | | 74- | 75- | 76- | 70- | 71- | 76- | 76- | 76- | |
| | | 14- | 32- | 00- | 21- | 00- | 14- | 21- | 21- | |
| | | 504C | 1662A | 118B | 528A | 24C | 396A | 833B | 702C | |
| No | Cult/Line | RPQQ | RTQS | RTQQ | QFBS | QSHS | TNMH | TNMK | RKQS | |
| 1 | MARQUIS | S | 23 | S | 2= | 2 | 2 | S | 23 | |
| 2 | CHRIS | 23 | 23 | 22- | 2 | ;1 | 0; | 2 | 2- | |
| 3 | Era | 0 . | 0 | . 0 | 0 | 2 | 0 | 0 | 0 | |
| 4 | STOA | _ | 2= | . 2= | 0; | 2= | 0 | 0 | 2= | |
| 5 | BUTTE 86 | 0 | 2= | 2= | 0; | 2= | 0 | 0 | 2= | |
| 6 | SD8072 | 2- | 2 | 2 | 2- | 23 | 2= | 2- | . 2 | |
| 7 | SD8073 | 2 | 23 | 2- | 2= | 2 | 2= | 2= | 2 | |
| 8 | SD8070 | 2= | 2 | 2= | 2= | 2- | 2= | 2= | 2 | |
| 9 | SD0005 | 0 | 0; | - | _ | - | - | ;1- | 0; | |
| 10 | SD0010 | 2- | ;1 | 2- | 0; | 23 | 0; | 2= | 0; | |
| 11 | MN89103 | 2= | 2 | 2= | 2= | 2= | ;1 | 2= | 2- | |
| 12 | MN90071 | 0 | ;1 | 1 | 0; | 2= | 0; | 0 | ;1 | |
| 13 | MN90114 | . 0 | 0; | 0 | 0 | 2= | 0 | 0 | 0; | |
| 14 | MN90253 | 0 | ;1- | 0 | 0 | 2= | 0 | 0 | 0; | |
| 15 | SBE0437 | 0;. | 0; | 0; | 0; | 2- | 0 | ;1- | 0; | |
| 16 | SBE0444 | 0; | ;1- | 0 | 0; | 1 | 0 | ;1- | ;1- | |
| 17 | ND671 | 0 | 2= | 2= | 0 | 2= | 0 | 0; | 2= | |
| 18 | ND673 | 2= | 2= | 2= | 2= | 2= | 2= | 2= | 2= | |
| 19 | ND674 | 2= | 2= | 2- | 2= | 2= | 2= | 2= | 2= | |
| 20 | ND677 | - | 2= | 2= . | 0; | 2= | 0 | 0; | 2= | |
| 21 | ND678 | - | 2= | 2= | 0 | 2= | - | 0 | 2= | |
| 22 | XW398A4 | 2= | 2= | 0 | 0; | 2= | 0 | - | 71- | |
| 23 | N86-0348 | _ | 2= | 0 | 0. | 2= | 0 | 0 | 2= | ; |
| 24 | N90-0666 | 0 | 0; | 0 | 0; | 2= | 0 | . 0 | 0; | |
| 25 | N90-0671 | 0 | ;1- | 0 | 0; | 1 | 0 | 0; | 0; | |
| 26 | N90-0700 | 2- | 23 | 2 | 2= | 2= | 2= | 2 | 2 | |
| | N88-3140 | – . | x | 2= | 0; | ;1 | 0; | 0; | 2= | |
| 8.5 | MT8849 | 0 | 2- | 0 | 2- | 2 - | 2= | 2- | 2- | |
| 29 | BW152 | ;1 | 2= | 0; | 0; | 0 | 0 | ;1 | 0; | |
| | 8601AE3C | ó | 23 | 2= | o · | 2= | 0 | Ó | 2- | |
| | BZ 988-351 | 2= | 0; | 2= | 0 | _ | 0 | 0 | 2= | |
| | BZ 984-334 | 0; | 0;,2 | = 2= | ;1 | 0,2= | = 0 | . 0 | 2= | |

Sr genes

| Set | I | 5 | 9d | 9e | 7b |
|-----|-----|----|----|----|-----|
| Set | II | 11 | 6 | 8 | 9a |
| Set | III | 36 | 9b | 13 | 10 |
| Set | IV | 15 | 16 | 17 | Tmp |

Table 4-1. Seedling reaction of the 1986 Hard Red Spring Wheat Performance Nursery to selected isolates of <u>Puccinia graminis</u> f. sp. <u>tritici</u>. (by D.V. McVey, USDA, ARS, U. of MN., St. Paul, MN).

| • | | | | | y Isol | | | |
|---------------------------|------------|------|-------|------------|--------|------|------------|--------------------|
| | 70- | 72- | 69- | 71- | 72- | 72- | 72- | 74- |
| • | 44- | 00- | 21- | 21- | 25 | 00- | 01- | 21- |
| | <u>68A</u> | 1370 | C 399 | 584B | 639C | 53A | 4A | 1490A |
| | 29 |] | 51 | 11- | -32-11 | .3 | 15 | B-2 |
| No. Entry | HJCS | QFBS | QSHS | RHRS | RKQS | RTQQ | TNME | TNMK |
| 1. Marquis | S | 2- | 2 | S | 23 | 23 | S | S |
| 2. Chris | 0 | 2- | X | 0; | 2 | 2 | 2 | 2 |
| Waldron | 2= | 2= | 2= | 2= | 2= | 2= | 2= | 2= |
| 4. Era | 0 | • | 2. | ; | ; | ; | ; | ; |
| 5. Butte | ; | ; | 2 | 2= | 2- | 2= | ; | ; |
| 6. SD2956 | 2- | 2= | 23cn | 23cn | 2= | 2= | 2= | |
| 7. SD2962 | , 0 | ; | 2= | ; | 2= | 2= | ; 1 | 2= |
| 8. SD2980(5 ha | .rp)0 | ; | 2-cn | 2-cn | 2= | 2= | <u>[j]</u> | $\exists \bigcirc$ |
| 9. SD2990 | ; | ; | 2 | ; I | 2- | 2= | ; | - ; |
| lO. SD2961 | ; | ; | 2- | ; | 2= | 2= | ; | ; |
| 1. MN82047 | ; | ; | 2 | 5 . | 1 | ; | ; | 2 |
| 12. MN81110 | 0 | ; | 2- | 3 - | ;1- | ; 1 | ; | ; |
| 13. MN82354 | ţ | ; | 2 | ; | 2 | 2 | ; | ; |
| 14. MN82008 | 0 | ; | lcn | 0; | ; | ; | ; | 2cn |
| 5. MN84662 | 0 . | ; | 2 | 0; | ; 1- | ; | ; | . ; |
| 6. MT8320 | ;1 | 2- | 2= | 2= | 23 | ; 1 | ;1 | 23 |
| 7. MT8304 | .0 | ; | 2 | 0; | 2= | ; | ; | ; |
| .8. MT8402 | ; . | ; | S | ; | 2 | ; | ; | ,1 |
| .9. ND606 | 2= | 2- | 2 | 2= | 2= | 21 | ; , | 2 ;,2 |
| 0. ND617 | ;1 | ; | ; | ; | 2= | 2= | 2= | 2= |
| 1. ND618 | 0 | ; | ; | ; | 2= | 2- | ; | ; |
| 2. ND626 | 2= | 2= | 2= | 2= | 2= | 2= | 2= | |
| 3. ND622 | ; | 1 | 2= | ; | 1- | 2= | 2= | 2= |
| 4. HS82-175 | 1 | 1 | 1 | 21 | 2 | 2= | 2= | 2= |
| 5. HS82-288 | 2= | 2= | 2= | 2 | 2- | 2= | 2= | 2- |
| 6. HS84-700 | 0 | ; | 2= | ; | ; | ; | ; | 2- |
| 7. HS84-873 | 2= | ; 1- | 2- | ; | 2= | 2- | 2= | 2- |
| 8. C982-324 | 2= | i | 2 | 2= | 2= | 2 | 2= | 2- |
| 9. WA7075 | 2= | 0; | 2 | ; | 2= | 2 | 2 | ; |
| 0. WA7329 | 0 | 0; | 1 | ; 1 | 2= | 2- | S | , , |
| 1. WA7330 | 0 | 2 | 2 | 32c | 2 | ; | • | 2 |

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE DIVISION BELTSVILLE, MARYLAND 20705

OBJECTIVE DESCRIPTION OF VARIETY WHEAT (Triticum spp.)

| NAME OF APPLICANT(S) | | |
|--|--|--|
| SOUTH DAKOTA AGRICULTURAL EXPERIMENT STATION | FOR OFFI | CIAL USE ONLY |
| DDRESS (Street and No. or R.F.D. No., City, State, and Zip Code) | PVPO NUMBER | 9600184 |
| | VARIETY NAME | |
| SOUTH DAKOTA STATE UNIVERSITY AG HALL 129 | | |
| BROOKINGS, SD 57007 | | |
| 3,00, | TEMPORARY OR E | CPERIMENTAL |
| | DESIGNATION | |
| | | en e |
| LEASE READ ALL INSTRUCTIONS CAREFULLY: Place the appropriate number that describes the varietal ace a zero in the first box (e.g or) when number is either 99 or less or 9 or less respectively. It is a minimum of 100 plants. Comparative data should be determined from varieties entered in the same trial. Resultant may be used to determine plant colors; designate system used: | yata for quantitative plant yal Horticultural Society | characters should be ba or any recognized color |
| 1=Common 2=Durum 3=Club 4=Other (SPECIFY VERNALIZATION: |) | |
| | | |
| 1=Spring 2=Winter 3=Other (SPECIFY) | ga kawara | |
| COLEOPTILE ANTHOCYANIN: | - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 | and the second of the second o |
| | | |
| 1=Absent 2=Present | | |
| JUVENILE PLANT GROWTH: | ·• | \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ |
| 2 1=Prostrate 2=Semi-erect 3=Erect | | |
| PLANT COLOR (boot stage): | | |
| 1 = Yellow-Green 2 = Green 3 = Blue-Green | | |
| FLAG LEAF (boot stage): | • 1 | |
| 2 1 = Erect 2 = Recurved EAR EMERGENCE: | 1 = Not Twisted | 2 = Twisted |
| ration that the control of the contr | ************************************** | |
| 0 3 Number of Days Earlier Than Chris | | |
| O 2 Number of Days Later Than Butte 86 | | |
| ANTHER COLOR: | | |
| 1 = YELLOW 2 = PURPLE | | |
| LANT HEIGHT (from soil to top of head, excluding awns): | Section 2 | |
| 0 5 cm Taller Than 2375 | e jakiya e | * |
| | | |

3 = Width Nearly as Wide as Kernel

2 = Depth 35% or less of Kernel

3 = Depth 50% or less of Kernel

| 960018 | | | | 9600184 |
|----------------|---|------------------------------|--|---------------------------------------|
| 13. SEED: (con | | - | | Exlubit C (Wheat) I |
| E. COI | I = White 2 = Amber | 3 = Red | 4 = Other (SPECIFY) | |
| F. TEX | TURE 1=Hard 2=Soft | | | |
| G. PHE | NOL REACTION (see instruction in the second | tions): 3 = Light Bro | own 4 = Dark Brown 5 = Black | |
| 14. DISEASE: | (0=Not Tested; 1=Suscer PLEASE IND | otible; 2=Res CATE THE SP | istant; 3=Intermediate; 4=Tolerant) ECIFIC RACE OR STRAIN TESTED | |
| | st <i>(Puccinia graminis</i> f. sp. <i>tritic</i> See Table 3 | 9 | Leaf Rust (Puccinia recondita f. sp. tritici) 2field reaction | |
| Stripe Ru | st (Puccinia striiformis) | 7 - | Loose Smut (Ustilago tritici) | |
| Tan Spot | (Pyrenophora tritici-repentis) | | Flag Smut (Urocystis agropyri) | 10001 1416-14 18001 14101 |
| Halo Spot | (Selenophoma donacis) | | Common Bunt (Tilletia tritici or T. laevis) | |
| Septoria n | odorum (Glume Blotch) | join . | Dwarf Bunt (Tilletia controversa) | |
| Septoria a | venae (Speckled Leaf Disease) | | Karnal Bunt (Tilletia indica) | |
| Septoria tra | itici (Speckled Leaf Blotch) | | Powdery Mildew (Erysiphe graminis f. sp. ti | itici) |
| Scab (Fus | arium spp.) | | "Snow Molds" | |
| | nt" (Kernel Smudge) | · · | Common Root Rot (Fusarium, Cochliobolu. | - :: |
| Barley Yell | ow Dwarf Virus (BYDV) | | Rhizoctonia Root Rot (Rhizoctonia solani) | |
| | Mosaic Virus (SBMV) | | Black Chaff (Xanthomonas campestris pv. tr | |
| | ow (Spindle Streak) Mosaic Vi | | Bacterial Leaf Blight (Pseudomonas syringa | |
| Wheat Stre | ak Mosaic Virus (WSMV) | | Other (SPECIFY) | |
| Other (SPI | ECIFY) | | Other (SPECIFY) | · · · · · · · · · · · · · · · · · · · |
| Other (SPE | ECIFY) | | Other (SPECIFY) | |
| Other (SPE | CCIFY) | | Other (SPECIFY) | |

| 5. INSECT: | (0=Not Tested; | 1=Susceptible; | 2=Resistant; | 3=Intermediate; | 4=Tolerant) | Exhibit C (Who |
|------------|---|-------------------------|--------------|---------------------------|-------------|-------------------------|
| | | PLEASE | SPECIFY BIOT | YPE (where needed) | | DENTRUM |
| Hessian F | ly <i>(Mayetiola destr</i> Heterogeneous | uctor) to Great Pla: | ins biotype | her (SPECIFY) | USI | RECEIVED DA-AMS-PVPO |
| Stem Saw | fly <i>(Cephus</i> spp.) | · | Ot | der (SPECIFY) | ' 96 | MAR 15 A10:39 |
| Cereal Le | af Beetle <i>(Oulema 1</i> | nelanopa) | Ott | ner (SPECIFY) | | |
| Russian A | phid <i>(Diuraphis no</i> | xia) Aleber | Oth | er (SPECIFY) | | |
| Greenbug 0 | (Schizaphis gramin | um) | Oth | er (SPECIFY) | | |
| Aphids | | egyenedî e. Politik | Oth | er (SPECIFY) | | |
| SeedI | ing tests at | the Cereal Ru | ıst Lab St | Paul, MN, indand/or Sr24. | | at the probable |

EXHIBIT D. Russ (SD8073) Additional Description of the Variety

The following additional descriptive information is presented:

- Release notice of Russ
- Table 4. South Dakota performance data.
- Table 5. Uniform Regional Hard Red Spring Wheat Nursery performance data.
- Table 6. 1994 Hard Spring Wheat Technical Committee data.

College of Agriculture and

Agricultural Experiment Station

Brookings, SD 57007-0291 Phone 605-688-4149 FAX 605-688-6065

Biological Sciences

Office of the Director Ag Hall 129, Box 2207

SDSU



DATE:

March 15, 1995

TO:

Agricultural Experiment Station Directors North Central Region and Great Plains

FROM:

Fred A. Cholick, Director

Agricultural Experiment Station

SUBJECT:

Release of Cultivars

The South Dakota Agricultural Experiment Station announces the release of the following cultivars and germplasms:

SD8073 Hard Red Spring Wheat - Proposed name 'Russ'; PVP recommended

21 SD8070 Hard Red Spring Wheat Germplasm

3) Spring Wheat Germplasm: SD-SWGP1, SD-SWGP2, SD-SWGP3, SD-SWGP4, and SD-SWGP5

The release date is March 1, 1995.

The South Dakota Agricultural Experiment Station also announces the increase with intent to release in 1996 SD0010 Hard Red Spring Wheat.

Enclosed are descriptions of the new releases and increase. Complete yield and agronomic data are available on request from the Plant Science Department.

Seed is available through the Foundation Seed Stocks Division, Department of Plant Science, South Dakota State University, Brookings, SD 57007 5418).

Action has also been taken to increase with intent to release the following crop varieties. Release date of the following materials will be dependent upon availability of seed. Information on this material is available on request from the Plant Science Department.

> SD302 Canada milk-vetch SD201 yellow-flowered alfalfa SD501 foxtail dalea SD92-1323M, SD92-1357M and SD92-1233M soybean lines SD89210, SD89504, and SD89210 oats

ph

Enclosures

Cultivar Release/Variety Recommendation Committee Plant Breeders



Northern Plains Biostress Laboratory Dedicated on September 17, 1993.

Creating Opportunities for a Lifetime

Educational programs and materials offered without regard to age, race, color, religion, sex, handicap or national origin. South Dakota State University, US Department of agriculture, and county cooperating. An Equal Opportunity Employer.

Release of SD8073 Hard Red Spring Wheat as 'Russ'

SD8073 hard red spring wheat will be released as 'Russ'. It is anticipated that Russ will be submitted for cultivar protection under the Plant Variety Protection Act, and Title V of the Federal Seed Act.

Russ is an awned, early maturing, standard height, hard red spring wheat. It is an F_6 derived line from the cross SD8052/SD2971. The pedigree of SD8052 is ND585/Shield and the pedigree of SD2971 is Agt/3/ND441//Wld/BB/4/Butte/5/Len. Russ has been in South Dakota State University trials since 1990 and in the Uniform Regional Hard Red Spring Wheat Nursery (HRSWURN) from 1991 to 1993. It was in the Spring Wheat Quality Advisory Council (SWQAC) trial in 1993 and 1994.

In relative performance to 'Butte 86' (109 site-years), SD8073 was 2 bushels/acre higher yielding, slightly lower in bushel weight, and similar in height. It is 2 days later heading than Butte 86 and 2 days earlier heading than 'Prospect'. South Dakota trials indicate that SD8073 yields more than Butte 86, Prospect, and 'Sharp'; and similar to '2375'. It appears to be widely adapted and has performed well in all areas tested in South Dakota as well as in HRSWURN locations.

Russ has good resistance to the prevalent races of leaf and stem rust in South Dakota. The probable genes for stem rust resistance are Waldron and/or SR24. It is heterogeneous for Hessian Fly resistance (H18), with approximately 50% of the plants resistant to the Great Plains biotype. Based on 1993 and 1994 performance data, Russ is similar to Butte 86 for Fusarium Head Blight resistance.

Russ is a strong mixing wheat with medium protein content. Three year quality evaluations by the U.S.D.A. Spring and Durum Wheat Quality Laboratory, Fargo, North Dakota, indicate that Russ is about 0.5% higher protein than Prospect and 0.2% lower protein than Butte 86; it has a longer mix time and a stronger mix pattern than Butte 86. In the SWQAC trials the protein content of Russ was between 0.5% and 1.0% less than 'Grandin', the milling characteristics were rated slightly lower than Grandin, and the baking properties were rated slightly higher than Grandin.

Table 4. Yield trial data from South Dakota.

SDSU Spring Wheat Breeding Trials 1990-1994 Grain Yield (bu/a)

| Ачеляде | (42) | 43.4 | 42.2 | 43.7 | 39.6 | 41.2 | | Ht | (30) | | 35 | 35 | 33 | 32 | 35 |
|-----------|---------------------------|--------|----------|------|----------|-------|---|-------|------|-----|--------|----------|------|----------|-------|
| Groton | 3 | 50.4 | 42.7 | 39.4 | 44.0 | 44.0 | | | (30) | | | | +2 | | |
| Day Co. | ଚ | 49.8 | 50.0 | 50.9 | 47.4 | 48.3 | | Prot | (18) | 199 | 13.9 | 14.1 | 13.7 | 13.3 | 13.0 |
| Brookings | () | 44.1 | 41.9 | 45.5 | 38.4 | 40.6 | | TW | (30) | | 57.3 | 58.1 | 59.0 | 56.3 | 59.2 |
| Raiph | ව | 39.8 | 40.7 | 41.5 | 41.2 | 38.3 | • | 90-94 | (42) | | 43.4 | 42.4 | 43.7 | 39.6 | 4.1 2 |
| Sclby | ଚ | 54.0 | 51.5 | 54.4 | 54.6 | 49.8 | | | | | | | | | |
| Highmore | 4 | 40.0 | 39.3 | 39.0 | 36.3 | 35.9 | | 1994 | (8) | | 42.3 | 38.9 | 38.3 | 37.8 | 40.2 |
| Brentford | € | 42.9 | 40.0 | 40.2 | 38.9 | 41.1 | | 1993 | 8 | | 38.4 | 37.7 | 39.7 | 34.6 | 36.2 |
| Watertown | ତ | 46.9 | 44.1 | 46.3 | 36.9 | 44.6 | | 1992 | 6) | | 57.5 | 54.3 | 58.5 | 52.0 | 53.3 |
| Redicid | (5) | 30.6 | 28.5 | 32.7 | 26.6 | 28.8 | | 1991 | 6) | | 33.4 | 35.2 | 35.9 | 31.2 | 33.0 |
| Brookings | ව | 40.3 | 41.6 | 43.9 | 36.1 | 40.5 | | 1990 | (8) | | 44.9 | 44.8 | 45.8 | 42.6 | 41.7 |
| | | SD8073 | Butte 86 | 2375 | Prospect | Sharp | | | | | SD8073 | Butte 86 | 2375 | Prospect | Sharn |

Crop Performance Trial, 1991-1994

| • | 1 | 1 | | G | rain Yie | snq) pra | shels/ac | re) | | | | ΛL | Hd | Hţ |
|----------|------------------|---|-----------------|-----------------|--------------|---------------|---|------|--------------|--------------|-------|--------|------|--------|
| | Brookings (4) | Watertown (4) | Highmore (3) | Redfield (4) | Selby (4) | Groton (4) | Selby Groton Aurora Co Wall (4) (4) (1) (4) | Wall | Bison (4) | Ralph (4) | 91-94 | lbs/bu | days | inches |
| SD8073 | 45.8 | 39.3 | 39.7 | 31.1 | 50.9 | 50.6 | 26.4 | 47.1 | 42.7 | 42.6 | 41.2 | 56.9 | +2 | 32 |
| Butte 86 | 44.3 | 36.3 | 39.1 | 27.3 | 50.2 | 51.7 | 23.4 | 47.1 | 37.2 | 43.2 | 39.9 | 58.6 | 0 | 33 |
| 2375 | 50.4 | 40.6 | 38.2 | 29.6 | 50.6 | 52.7 | 33.9 | 45.4 | 39.0 | 43.8 | 41.4 | 58.5 | +5 | 31 |
| Prospect | 45.5 | 34.2 | 39.5 | 27.3 | 51.0 | 52.3 | 29.1 | 45.2 | 42.0 | 47.5 | 40.7 | 56.6 | +3 | 30 |
| Sharp | 48.2 | 35.8 | 38.0 | 28.1 | 47.4 | 52.7 | 28.5 | 45.6 | 37.8 | 40.1 | 40.0 | 57.6 | +1 | 33 |
| - | | | | | | | | | | | | | | |

Table 5. Yield trial data from the Uniform Regional Hard Red Spring Wheat Nursery. 1991-1993: North Dakota, South Dakota, Minnesota.

| | 1991 (11) | 1992 (10) | 1993 (10) | 91-93 (31) | |
|----------|--------------|--------------|---------------|---------------|---|
| | | Grain Yi | eld (bu/acre) | | |
| Stoa | 35.3 | 55.4 | 36.3 | 42.3 | |
| Butte 86 | 37.2 | 52.1 | 37.6 | 42.3 | • |
| Russ | 38.8 | 57.9 | 41.7 | 46.2 | |
| Kulm | 38.7 | 52.1 | 39.7 | 43.5 | |

Characteristics (1993)

| 44 | TW Heading lb/bu days | | Height (cm) | Prot % | Stem Rust Leaf Rust | | |
|----------|--------------------------|------|-------------|-----------|---------------------|---------|--|
| | (16) | (15) | (16) | (13) | (1) | (1) | |
| Stoa | 56.0 | 34 | 100 | 13.9 | 0 | 10MR-MS | |
| Butte 86 | 57.4 | 30 | 92 | 13.8 | 5RMR | 10MR-MS | |
| Russ | 56.1 | 32 | 92 | 13.4 | TMR | TMR-MS | |
| Kulm | 58.3 | 30 | 96 | 13.9 | TR | TR-MR | |

Leaf rust and stem rust are from an inoculated nursery in St. Paul.

Table 6. 1994 HARD SPRING WHEAT TECHNICAL COMMITTEE

VARIETY: SD8073

| | | | | | | | | | Average | Average |
|---------------------------|------|------|------|------|------|------|------|--------------|---------|---------|
| Sample Code: | | | C-CK | | K-CK | K-12 | M-CK | M-12 | Grandin | SD8073 |
| Wht Protein(14%mb): | | 15.1 | 14.5 | 13.7 | 14.5 | 13.6 | 16.1 | 15.2 | 15.3 | 14.4 |
| Wheat Ash(14%mb): | 1.72 | 1.74 | 1.67 | 1.79 | 1.69 | 1.66 | 1.55 | 1.47 | 1.66 | 1.67 |
| Test Weight(lb/bu): | | 57.1 | 59.8 | 59.7 | 60.4 | 60.5 | 62.0 | 62. 6 | 59.9 | 60.0 |
| 1000-KWT(gram): | | 28.5 | 31.6 | 31.7 | 30.7 | 30.8 | 36.4 | 35.8 | 31.7 | 31.7 |
| Large Kemels(%): | 49 | 46 | 54 | 52 | 58 | 62 | 69 | 74 | 58 | 59 |
| Small Kemels(%): | | 1 | 2 | 2 | 2 | 1 | 1 | 1 | 2 | 1 . |
| NIR Hard: | 90.0 | 81.0 | 95.0 | 87.0 | 97.0 | 97.0 | 90.0 | 94.0 | 93 | 90 |
| Kernel Vit(%): | 60.9 | 44.2 | 75.6 | 74.0 | 80.3 | 85.7 | 93.3 | 71.1 | 77.5 | 68.8 |
| SKWCS HI: | 78.5 | 74.9 | 81.4 | 84.0 | 85.2 | 86.8 | 70.2 | 71.8 | 78.8 | 79.4 |
| Wht FN(sec): | 352 | 385 | 384 | 388 | 371 | 377 | 363 | 366 | 368 | 379 |
| FI Protein(14%mb): | 15.2 | 14.0 | 13.7 | 13.1 | 13.9 | 13.1 | 15.3 | 14.3 | 14.5 | 13.6 |
| Fl Ash(14%mb): | 0.46 | 0.43 | 0.43 | 0.49 | 0.47 | 0.46 | 0.41 | 0.40 | 0.44 | 0.44 |
| FI Ext(%): | 70.7 | 68.9 | 72.6 | 71.5 | 73.5 | 70.5 | 72.3 | 72.1 | 72.3 | 70.7 |
| # .46 Ash Fl/cwt Wht: | 73.9 | 72.5 | 74.6 | 62.2 | 73.5 | 71.9 | 76.9 | 76.8 | 74.7 | 70.9 |
| Mill Value(\$): | 1.89 | 1.79 | 1.91 | 1.68 | 1.93 | 1.80 | 2.07 | 2.11 | 1.95 | 1.85 |
| Farino Abs(14%mb): | 62.4 | 62.2 | 61.1 | 61.7 | 60.5 | 60.5 | 66.1 | 64.6 | 62.5 | 62.3 |
| Farino Arrival Time(min): | 3.0 | 2.7 | 2.8 | 2.3 | 2.3 | 1.9 | 4.2 | 3.9 | 3.1 | 2.7 |
| Farino Peak Time(min): | 6.5 | 9.1 | 5.2 | 5.5 | 4.5 | 5.9 | 7.6 | 6.4 | 6.0 | 6.7 |
| Farino Stability(min): | 11.9 | 15.3 | 8.1 | 9.8 | 8.1 | 10.3 | 12.4 | 9.6 | 10.1 | 11.3 |
| Farino MTI(BU): | 14 | 24 | 29 | 22 | 28 | 30 | 9 | 19 | . 20 | 24 |
| Bake Asorption(14% mb): | 62.4 | 62.2 | 60.3 | 60.9 | 60.4 | 60.6 | 64.1 | 64.0 | 61.8 | 61.9 |
| Bake Rating: | 3.8 | 3.4 | 2.8 | 2.9 | 2.8 | 2.9 | 4.5 | 4.4 | 3.5 | 3.4 |
| Bake Mix Time Actual: | 11.0 | 11.4 | 8.5 | 8.8 | 8.2 | 9.1 | 11.5 | 10.1 | 9.8 | 9.8 |
| Bake Mix Time Rating: | 3.9 | 4.3 | 3.0 | 3.0 | 2.5 | 3.3 | 4.2 | 3.7 | 3.4 | 3.6 |
| Mix Tolerance Rating: | 3.8 | 4.0 | 2.5 | 3.3 | 2.7 | 3.1 | 4.4 | 3.7 | 3.4 | 3.5 |
| Out of Mixer Rating: | 3.6 | 3.7 | 3.3 | 3.3 | 2.9 | 3.1 | 3.9 | 3.5 | 3.4 | 3.4 |
| Out of Mixer Describe: | 2.1 | 2.3 | 1.8 | 2.0 | 1.8 | 2.2 | 2.3 | 2.1 | 2.0 | 2.1 |
| At Make Up Rating: | 3.9 | 3.7 | 2.8 | 3.3 | 2.5 | 3.6 | 4.3 | 3.4 | 3.4 | 3.5 |
| At Make Up Describe: | 2.2 | 2.3 | 1.5 | 2.1 | 1.5 | 1.8 | 2.5 | 2.4 | 1.9 | 2.2 |
| Loaf Volume Rating: | 3.8 | 3.9 | 3.4 | 3.2 | 3.0 | 3.8 | 4.8 | 3.7 | 3.7 | 3.6 |
| Crumb Color: | 2.7 | 3.8 | 2.9 | 3.8 | 3.1 | 4.0 | 3.8 | 4.0 | 3.1 | 3.9 |
| Crumb Grain: | 3.4 | 3.8 | 3.2 | 3.4 | 3.5 | 4.1 | 3.9 | 3.8 | 3.5 | 3.8 |
| Crumb Texture: | 3.5 | 3.6 | 2.9 | 3.8 | 3.4 | 4.0 | 3.8 | 3.8 | 3.4 | 3.8 |
| Overall Rating: | 3.3 | 3.7 | 2.7 | 3.3 | 3.0 | 3.6 | 4.0 | 3.7 | 3.2 | 3.6 |

| Rating Scores: 0 | 3 | 6 |
|---------------------------|-----------------------|------------------------|
| Bake Absorption: Low | | High |
| Bake Mix Time: Short | | Long |
| Mixing Tolerance: Weak | | Strong |
| Out Of Mixer: Weak | · | Bucky |
| At Make Up: Weak | | Bucky |
| Loaf Volume : Low | | High |
| Crumb Color: Yellow | Grey Dull Creamy | Bright White |
| Crumb Grain : Irregular,o | pen,thick Open, thick | Close, elongated, fine |
| Crumb Texture: Harsh | | Silky |
| Overall Rating: Poor | | Excellent |

Out of Mixer Describe:

- 1. Sticky Weak
- 2. Medium Pliable
- 3. Tough Bucky

Out of Mixer Describe:

- 1. Sticky Weak
- 2. Medium Pliable
- 3. Tough Bucky

| REPRODUCE LOCALLY. Include form number and date on all reproductions. | FORM APPROVED - OMB N | 0. 0581-0055 EXPIRES: 12-31-9 |
|---|---|--|
| U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY DIVISION - PLANT VARIETY PROTECTION OFFICE EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP | 1974 (5 U.S.C. 552a) and the Pap Application is required in order to | e in accordance with the Privacy Act erwork Reduction Act (PRA) of 1995 . determine if a plant variety protection 2421). Information is held confidential |
| 1. NAME OF APPLICANT(S) | 2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER | 3. VARIETY NAME |
| SOUTH DAKOTA AGRICULTURAL EXPERIMENT STATION | SD 8073 | Russ |
| 4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code, and Country) | 5. TELEPHONE (include area code) | 6. FAX (include area code) |
| SOUTH DAKOTA STATE UNIVERSITY | 605-688-4149 | 605-688-6065 |
| AG HALL 129 BROOKINGS, SD 57007 | 7. PVPO NUMBER 9600184 | • |
| 8. Does the applicant own all rights to the variety? Mark an "X" in appropriate b | vilock. If no, please explain. | X YES NO |
| | | |
| Is the applicant (individual or company) a U.S. national or U.S. based company If no, give name of country | ? | X YES NO |
| a. If original rights to variety were owned by individual(s): Is (are) the original breeder(s) a U.S. national(s)? If no, give name of c | ountry | YES NO |
| b. If original rights to variety were owned by a company: ls the original breeder(s) U.S. based company? If no, give name of course. | entry | |
| 11. Additional explantion on ownership (If needed, use reverse for extra space): | | : |
| | | |
| PLEASE NOTE: | | |
| Plant variety protection can be afforded only to owners (not licensees) who meet or | ne of the following criteria: | |
| f. If the rights to the variety are owned by the original breeder, that person must be of a country which affords similar protection to nationals of the U.S. for the same | pe a U.S. national, national of a U ne genus and species. | POV member country, or national |
| If the rights to the variety are owned by the company which employed the origin nationals of a UPOV member country, or owned by nationals of a country which genus and species. | nal breeder(s), the company must affords similar protection to natio | be U.S. based, owned by nals of the U.S. for the same |
| . If the applicant is an owner who is not the original breeder, both the original bree | eder and the applicant must meet | one of the above criteria. |
| he original breeder may be the individual or company who directed final breeding efinition. | ng. See Section 41(a)(2) of the | Plant Variety Protection Act for |
| | | |

Public reporting burden for this collection of information is estimated to average 10 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Officer, OIRM, AG Box 7630, Jamie L. Whitten Building, Washington, D.C. 20250. When replying, refer to OMB No. 0581-0055 and form number in your letter.

Under the PRA of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA Office of Communications at (202) 720-2791.

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call (202) 720-7327 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.